

CONCISE DEFINITIVE REVIEW

Bram Rochwerf, MD, MSc (Epi), FRCPC, Series Editor

Assessment of Tissue Perfusion Pressure in Patients With Septic Shock: Beyond Mean Arterial Pressure

OBJECTIVES: Optimization of macrohemodynamics is just the starting point in the management of hemodynamics in patients with septic shock. We describe the interaction between the various determinants of arterial pressure and tissue perfusion, how to optimize them and their estimations at the bedside. This is a concise definitive review of the assessment of tissue perfusion pressure (TPP) in patients with septic shock beyond just mean arterial pressure (MAP).

DATA SOURCES: Original publications were retrieved through a PubMed and MEDLINE databases with search terms related to septic shock, arterial blood pressure, critical closing pressure (Pcc), mean systemic filling pressure (Pmsf), and TPP. Supporting evidence was also retrieved from PubMed and MEDLINE when indicated.

STUDY SELECTION: English-language systematic reviews, narrative reviews, meta-analyses, randomized clinical trials, and observational studies.

DATA EXTRACTION AND DATA SYNTHESIS: Data from relevant publications were reviewed, selected, and summarized by the authors and applied as indicated.

CONCLUSIONS: The relationship between arterial pressure and outcomes in septic shock is complex and heterogeneous. Focusing on critical parameters—such as TPP, the vascular waterfall, and individual treatment effects—enables a more personalized approach, focused on perfusion rather than pressure alone. Persistent hypoperfusion despite adequate macrocirculatory targets marks a pivotal moment when microcirculatory assessment becomes essential to guide therapy and avoid potentially harmful escalation of fluids or vasopressors. Bedside measurements of systolic arterial pressure, MAP, diastolic arterial pressure, Pcc, and Pmsf offer practical tools for monitoring and tailoring treatment. Future clinical trials are needed to validate these metrics and refine resuscitation strategies in septic shock.

KEYWORDS: arterial blood pressure; critical closing pressure; mean systemic filling pressure; septic shock; tissue perfusion pressure

Carlos Sanchez E. , MD¹Ahmed Taha, MD²Yasser Tolba, PhD^{3,4}Glenn Hernandez, PhD⁵Michael R. Pinsky, MD⁶

Septic shock is a life-threatening condition (1) associated with vasoplegia that impairs tissue perfusion, leading to organ dysfunction and death (2). Both the depth and duration of arterial hypotension impact mortality (3–5). Restoring mean arterial pressure (MAP) and cardiac output (CO) to threshold targets by fluids and vasopressors may not restore organ function or tissue perfusion (6–8). The dissociation between measurable macrocirculatory parameters and microcirculatory flow warrants deeper analysis of arterial pressure and tissue blood flow determinants.

Copyright © 2025 by the Society of Critical Care Medicine and Wolters Kluwer Health, Inc. All Rights Reserved.

DOI: 10.1097/CCM.0000000000006805





KEY POINTS

Question: Can we assess tissue perfusion pressure (TPP) and vasomotor tone in patients with septic shock beyond mean arterial pressure (MAP)?

Findings: MAP relative to critical closing pressure (Pcc) defines TPP, while the pressure gradient between Pcc and mean systemic filling pressure (Pmsf) defines the vascular waterfall essential to allow for autoregulation of local flow. Low vasomotor tone is characterized by a decrease in diastolic arterial pressure. Importantly, tachycardia, by limiting diastolic runoff time, minimizes hypotension, which is good, but masks vasoplegia. To the extent that hemodynamic factors in sepsis create inadequate tissue perfusion, resuscitation efforts that restore both TPP and local vascular waterfalls should improve the linkage between macrocirculatory parameters, such as MAP and cardiac output, with microcirculatory measures of tissue perfusion. This broader view of the determinants of septic shock-induced tissue hypoperfusion should allow for a more nuanced approach to the use of fluids, vasopressors, and inotropes.

Meaning: Arterial blood pressure measures and their derived tissue perfusion parameters may be estimated at the bedside and could potentially guide clinical decision-making beyond MAP in patients with septic shock.

ARTERIAL PRESSURE AND THE DETERMINANTS OF ORGAN BLOOD FLOW

Arterial pressure is not a single number nor solely defines pathology or organ perfusion. Understanding its determinants is crucial for clinical decision-making. The four main parameters defining the arterial profile are systolic arterial pressure (SAP), diastolic arterial pressure (DAP), pulse pressure (PP), and MAP (9) (**Fig. 1**). SAP is the peak pressure from left ventricular (LV) ejection, arterial tone, reflected waves, and arterial viscoelastic properties. DAP is the minimal pressure before ejection, determined by central blood volume, vasomotor tone, and heart rate (HR). PP is the difference between DAP and SAP, influenced by stroke volume (SV), compliance, and SAP/DAP




Figure 1. Components of arterial blood pressure. DAP = diastolic arterial pressure, MAP = mean arterial pressure, PP = pulse pressure (systolic arterial pressure–diastolic arterial pressure), SAP = systolic arterial pressure.

factors. MAP, is an arithmetic value of arterial pressure across a heartbeat, is calculated as DAP plus one-third of PP (10).

Consensus definitions of septic shock include hypotension (e.g., $SAP \leq 100$ mm Hg) or vasopressor need to maintain MAP (1). Guidelines recommend targeting an initial MAP greater than or equal to 65 mm Hg (11), a reasonable target as SAP varies across the arterial circuit (12–14). SAP site-specific differences may impair resuscitation if minimal SAP is the target. Low DAP reflects vasoplegia; MAP, unlike SAP, is relatively constant throughout the arterial tree and reflects average arterial pressure (10).

MAP constancy down to small arteries/arterioles reflects large vessels' role as capacitors maintaining pressure during diastole with little forward flow. As arteries branch into smaller vessels, the total cross-sectional area increases, vascular resistance rises, and blood pressure drops rapidly over a short distance (15). When intraluminal pressure falls below the vessel wall tone from smooth muscle, vessels collapse. This pressure defines the arterial critical closing pressure (Pcc) (16, 17). The collapse point moves distally with vasomotor tone loss, as in sepsis, or more proximally with tone increase, as in early hemorrhagic or cardiogenic shock (16). Tissues regulate local blood flow by altering vasomotor tone in small feeding arteries/arterioles to meet metabolic demands (18).

Tissue perfusion pressure (TPP) is the gradient between MAP and Pcc. Blood flow changes occur via feedback-driven tone changes through endothelial retrograde signaling (16). For Pcc changes to affect blood flow, two conditions must be met: TPP must sustain forward flow (19) and Pcc must exceed downstream capillary pressure enough to drive flow (15, 20, 21). Capillaries' large cross-sectional area and slow flow mean minimal back pressure, except for venular pressure. Although Pcc varies by tissue, venular backpressure is relatively constant due to high capacitance and area of drainage vessels and their slow flow (22). This backpressure is called mean systemic filling pressure (Pmsf)—the pressure after abrupt flow cessation and redistribution, estimating upstream pressure driving venous return and correlating with circulatory fullness (23).

Normally, Pmsf is low (≤ 10 mm Hg), and Pcc is high (~ 45 mm Hg at rest), decreasing with increased metabolic demand. This creates a “vascular waterfall”

between arterial inflow (Pcc) and capillaries (24). Pmsf depends on venular tone, tissue pressure, and blood volume. Bedside methods exist to estimate Pmsf (25), and a smartphone app called iGuyton (26) estimates Pmsf and venous return parameters from demographic data, CO, MAP, and right atrial pressure (Pra). The venous return pressure gradient is Pmsf–Pra (10). **Figure 2** illustrates these concepts and their circulation control points.

The vascular waterfall protects capillaries from high pressure and edema. However, increased metabolic demand reduces Pcc and raises capillary pressure, promoting fluid translocation (27). During exercise, local Pcc drops, causing interstitial edema, “muscle bulking” reflects this. Although each tissue has specific Pcc values, a global Pcc estimate can be derived from systemic pressure changes during respiration or sinus pauses (28). MAP, SAP, DAP, Pcc, and Pmsf can be bedside-estimated with non- or minimally invasive





Figure 2. Didactic illustration of different pressures that can be measured at each control point along the circulatory system. CVP = central venous pressure, Pcc = critical closing pressure, Pmsf = mean systemic filling pressure, Pra = right atrial pressure, TPP = tissue perfusion pressure.

Septic shock induces generalized vasoplegia by blunting vascular endothelial-smooth muscle feedback (2). This decreases both Pcc and Pmsf, but Pcc decreases more due to its unique determinants. Clinically, Pcc may fall to levels near Pmsf (12). In such cases, auto-regulation is lost. Although resuscitation may increase MAP and CO, the resulting flow is uniformly distributed, leading to underperfusion of metabolically active tissues and overperfusion of inactive ones. This state is hyperdynamic hypotension: low MAP and DAP, high CO, HR, and mixed venous oxygen saturation, along with hyperlactatemia and organ dysfunction (2).

sepsis, systemic inflammation can trigger intracellular metabolic changes that persist despite adequate TPP and vascular waterfall restoration (29, 30). Clinicians may face patients in septic shock with normalized MAP and CO, yet persistent organ dysfunction. Measuring TPP, Pcc, and presence of vascular waterfall may guide further care: either intensifying resuscitation if inadequate or holding support if values are sufficient since more fluids or vasopressors might be harmful. However, this has not been tested in trials.

Ventriculo-Arterial Coupling

The LV ejects SV into a pressurized arterial system. SV generates forward flow, but arterial pressure limits output due to afterload. As impedance increases, SV falls depending on the balance between end-systolic



Sepsis reduces arterial elastance via vasoplegia (2, 32). Thus, for the same contractility and end-diastolic volume, SV increases while SAP stays constant or decreases. Sepsis may also impair contractility, but this is rarely apparent in early hyperdynamic septic shock, as low arterial elastance masks it. Increasing elastance with vasopressors can reveal poor contractility, seen as reduced CO (31, 33, 34).

Figure 3. Schematic presentation of ventricular-arterial coupling on the pressure-volume relationship. E_a = arterial elastance, EDP = end-diastolic pressure, EDPVR = end-diastolic pressure-volume relationship, EDV = end-diastolic volume, E_{es} = end-systolic elastance, EF = ejection fraction, ESP = end-systolic pressure, ESPVR = end-systolic pressure-volume relationship, ESV = end-systolic volume, PE = potential energy, SAP = systolic arterial pressure, SV = stroke volume, SW = stroke work, VAC = ventricular-arterial coupling, V_o = volume axis interception.

The free smartphone app iElastance (35) estimates end-systolic elastance, arterial elastance, and VAC using SAP, DAP, ejection fraction, SV, pre-ejection time, and ejection time.

Dynamic Arterial Elastance

Changes in LV SV induce proportional changes in PP. This relationship depends on central arterial compliance, termed dynamic arterial elastance (E_{adyn}) (36). E_{adyn} is calculated as the ratio of pulse pressure variation (PPV) to SV variation, measured during spontaneous or positive-pressure ventilation (31, 34). E_{adyn} is related to VAC (37) but primarily assesses how MAP will change with altered CO or vasopressor use (38, 39).

In hypotensive, volume-responsive patients, CO may rise with fluids, but MAP might not increase. This is common in septic shock due to low arterial elastance. If E_{adyn} is less than 1 before fluid loading, MAP typically increases little. These patients benefit from vasopressors to shorten hypotension duration and reduce fluid needs (34, 40).

Conversely, in vasoplegic patients on vasopressors, it's unclear if support is still needed. Currently, clinicians test weaning by lowering vasopressors and observing MAP. If it falls, the patient is vasopressor-dependent. However, this trial can trigger hypotension and worsen organ injury. If E_{adyn} is greater than 1.1, vasopressor weaning rarely causes hypotension, enabling safer, faster tapering in stable patients (38, 41–43).

CLINICAL RELEVANCE OF ARTERIAL PRESSURE COMPONENTS

Systolic Arterial Pressure

LV contraction propels blood into the aorta, raising central arterial pressure. This rise occurs during systolic ejection, making SAP a measure of LV performance. Arterial stiffness, or elastance, is a secondary determinant of SAP (15, 44). Additionally, backward-reflected pressure waves from vascular branching points further elevate peripheral SAP (pSAP) (45) (**Fig. 4**). Clinically, SAP is a quick triage marker, easily measured by sphygmomanometry, and commonly used to define hypotension in acute settings.

SAP less than or equal to 100 mm Hg is an early sign of sepsis (1), but it also appears in cardiogenic,

hypovolemic, and obstructive shock. Thus, systolic hypotension is a nonspecific marker of circulatory failure (46). Literature on SAP's prognostic value is mixed, but higher SAP often correlates with better outcomes (47). Marked, irregular SAP swings suggest higher mortality risk and are more likely related to fluid deficiency than vasopressor need (48).

Mean Arterial Pressure

MAP is the average arterial pressure during a cardiac cycle, estimated as DAP plus one-third the PP. Normal MAP ranges from 70 to 110 mm Hg (49). In septic shock, 65 mm Hg is considered the minimum initial threshold for end-organ perfusion (50) or no more than 15% below baseline in chronic hypertensive patients (51, 52). Since MAP represents only input pressure, high venous or intra-abdominal pressure may demand a higher MAP to maintain perfusion. It's the TPP, not MAP alone, that determines tissue flow (53, 54).

MAP reflects the overall cycle pressure and is heavily influenced by DAP, as the heart spends more time in diastole.

Diastolic Arterial Pressure

DAP is the lowest arterial pressure during the cardiac cycle. It's influenced by residual arterial volume at diastole's end, vessel tone, and Pcc. Diastolic time impacts this volume, and tachycardia shortens it, raising DAP for the same SV (55–57). Patients with sepsis and chronic hypertension may need a DAP of 54–62 mm Hg for adequate perfusion (52).

Coronary blood flow to the LV occurs during diastole, as systolic contraction compresses coronary vessels. Therefore, DAP is the input pressure for LV perfusion. If DAP falls below 40 mm Hg, myocardial ischemia may occur (58). DAP less than 40 mm Hg is often due to reduced vascular tone, especially in tachycardia (59, 60). Xiao et al (61) identified optimal DAP and HR ranges of 50–70 mm Hg and 60–90 beats/min (bpm) in septic shock. DAP less than or equal to 40 mm Hg with HR greater than or equal to 100 bpm predicts higher mortality.

Holder et al (62) showed DAP less than 52 mm Hg predicts early septic shock progression within 96 hours. Lower DAP indicates greater vasoplegia




Figure 4. Augmentation pressure phenomenon. **A**, Forward arterial pressure wave. **B**, Reflected pressure wave from arterial reflectance points coming retrograde from the periphery to the central circulation. **C**, Impact of wave reflection on the pressure wave contour increasing systolic blood pressure (augmentation pressure) due to earlier reflected wave in patients with increased arterial stiffness (*right side*). The *red line* shows the resulting arterial waveform. *Resulted normal and abnormal waveforms are directly related to how elastic or stiff (respectively) is the specific artery where blood pressure is being measured. P1 = ejection wave, P2 = reflecting wave, P3 = augmentation pressure.

and worse outcomes. Gao et al (63) found an inverse link between DAP and 28-day mortality post-resuscitation—DAP less than 59 mm Hg at 24 hours correlated with poor survival, suggesting targeting DAP greater than or equal to 59 mm Hg after achieving MAP 65 mm Hg may improve outcomes. This concept is being explored in the ongoing. A multicenter randomized clinical trial designed to compare the effects of peripheral perfusion-targeted resuscitation versus lactate-guided resuscitation on mortality in patients with septic shock (ANDROMEDA-SHOCK-2) trial (64).

Even if MAP exceeds 65 mm Hg, low DAP is still linked to high 90-day mortality (51, 65). Furthermore, microcirculatory dysfunction often persists after MAP

normalization (66, 67). Collet et al (68) identified DAP as the only independent determinant of tissue oxygen resaturation in sepsis.

Pulse Pressure

PP is the difference between SAP and DAP. It noninvasively reflects SV relative to vasomotor tone. In sepsis, low PP is tied to higher mortality (69–71). Respiratory-induced changes in PP, or PPV, are common (72). In ventilated patients, PPV greater than or equal to 13% predicts fluid responsiveness with high sensitivity (85–89%) and specificity (88–98%) (73).

Various respiratory maneuvers help evaluate fluid responsiveness using PPV (74). PPV is a simple

bedside tool to predict and assess the hemodynamic impact of intravascular volume expansion in septic patients.

INFLUENCE OF HEART RATE ON ARTERIAL BLOOD PRESSURE

HR exerts a significant influence on arterial blood pressure (ABP), particularly through its effect on diastolic time. Tachycardia shortens the diastolic phase, thereby limiting arterial runoff and resulting in elevated DAP for the same SV (10). This phenomenon can mask underlying vasoplegia, as the rise in DAP is not due to increased vascular tone but to the reduced time available for diastolic runoff (**Fig. 5**). Reflex tachycardia in critically ill patients may therefore give the false impression of preserved vascular tone.

Wilkinson et al (45) demonstrated that incremental pacing, which increases HR without affecting CO, results in a proportional decrease in SV but still leads to increased SAP, DAP, and MAP. Interestingly, while central DAP increased with pacing, central SAP (cSAP) did not show the same trend, likely due to differences in the timing of reflected pressure waves. This underscores the complex interplay between HR and the components of ABP.

A clinically relevant parameter that integrates HR and DAP is the diastolic shock index (DSI), defined as HR divided by DAP. Ospina-Tascón et al (75) found

that a DSI greater than 2.2 was associated with significantly higher mortality in patients with septic shock. The relationship is such that for a constant HR, a lower DAP increases mortality, and for a constant DAP, a higher HR has the same effect. Notably, both HR and DAP are independent predictors of mortality in this setting.

BEDSIDE ARTERIAL PRESSURE MONITORING

Although invasive ABP monitoring is frequently employed in critically ill patients, evidence does not uniformly support its routine use. This is largely because measurements alone, without being used to guide targeted interventions, do not necessarily improve clinical outcomes. Kaufmann et al (76) compared noninvasive and invasive measurements of SAP, MAP, and DAP in patients, including those on norepinephrine infusions, and found no significant differences between the two modalities. However, a meta-analysis by Kim et al (77) concluded that continuous noninvasive ABP monitors tend to be both inaccurate and imprecise beyond acceptable thresholds mainly when the patient is hypotensive (78). These discrepancies are especially relevant when calculating DSI or making critical decisions in hypotensive patients. As a result, invasive arterial monitoring remains the standard recommendation in unstable patients, according to major critical care guidelines.

In vasoconstricted states, peripheral ABP readings may become unreliable. Kim et al (79) showed that in patients receiving greater than or equal to 0.1 µg/kg/min of norepinephrine to maintain MAP greater than or equal to 65 mm Hg, radial artery pressures underestimated femoral artery pressures by greater than or equal to 5 mm Hg in 62.2% of cases. Wisanusattra and Khwannimit (80) reported similar findings. This discrepancy arises because pSAP is often higher than cSAP due to reflected pressure wave amplification (78, 81). Therefore, pSAP is not a reliable surrogate for assessing central hemodynamics or cardiovascular risk (82). To correct for this, Chemla et al (83) proposed an equation to estimate cSAP from peripheral MAP and DAP: $cSAP = MAP^2/DAP$. This approach minimizes reliance on potentially misleading pSAP values. Importantly, noninvasive methods tend to overestimate DAP by




Figure 5. Influence of the heart rate on arterial blood pressure. Black waves represent more or less constant values of systolic arterial pressure (SAP) and diastolic arterial pressure (DAP) at normal heart rates. The red waves show how as heart rate increases, arterial blood pressure also increases, with a greater increase in DAP than in SAP with respect to the base line values. MAP = mean arterial pressure.

an average of 6 mm Hg and underestimate SAP by the same amount when compared with invasive measures (84, 85). In vasoplegic states, reflected pressure waves are diminished, causing peripheral measurements to underestimate central pressures even more markedly (86).

TISSUE PERFUSION PRESSURE

TPP, defined as the difference between MAP and the Pcc, reflects the driving force for tissue blood flow. Since Pcc is typically higher than the Pmsf, local vasodilation (which decreases Pcc) can enhance perfusion. This is the primary mechanism by which the body matches local blood flow to metabolic demand through autoregulation.

However, in vasoplegia, Pcc may fall to levels approaching Pmsf, impairing this autoregulatory mechanism. In such cases, even increases in MAP may not improve perfusion. A TPP greater than 34 mm Hg has been associated with better lactate clearance and improved microcirculatory regulation in cardiac surgery patients (19). Andrei et al (87) observed that in

postoperative vasoplegic patients, the Pcc–Pmsf gradient was less than 5 mm Hg before resuscitation. Following norepinephrine administration to raise MAP above 80 mm Hg, only half of these patients showed an increase in Pcc, reestablishing a vascular waterfall (Pcc >> Pmsf). Those who did showed better perfusion outcomes, including improved lactate clearance, lower delta PCO₂, improved sublingual microcirculatory blood flow, and faster capillary refill time (CRT) (87).

PCC AND PMSF BEDSIDE MEASUREMENT

Multiple bedside techniques have been developed to estimate Pcc and Pmsf (Table 1).

Pcc

Each vascular bed has its own Pcc, which varies with metabolic demand. However, a general Pcc can be estimated bedside by rapidly occluding a peripheral arterial catheter (28); the initial arterial pressure decay

TABLE 1. Bedside Measurement of Critical Closing Pressure and Mean Systemic Filling Pressure

Pcc Bedside Measurement		
Name of Method	Technique	Type of Patient
Pcc	Peripheral arterial catheter measuring arterial pressure is rapidly occluded. The initial arterial pressure decay during the first 10–12 s (approximates Pcc)	All of them
Pmsf Bedside Measurement		
Name of Method	Technique	Type of Patient
Pmsf arm	Arm cuff is rapidly inflated stopping forearm flow (arterial and venous) equilibrium pressure corresponds to Pmsf	Awake or sedated Spontaneous breathing or on mechanical ventilation Sinus or cardiac arrhythmias
Pmsf analogue	Guytonian mathematical model using the following formula: Pmcf analogue = a × CVP + b × MAP + c × CO	Awake or sedated Spontaneous breathing or on mechanical ventilation Sinus or cardiac arrhythmias
Pmsf inspiratory hold	End-inspiratory hold maneuvers, simultaneously measure changes in right atrial pressure and cardiac output. The zero right atrial pressure interception value corresponds to Pmsf	Must be on mechanical ventilation with no spontaneous ventilatory efforts Without cardiac arrhythmias

CO = cardiac output, CVP = central venous pressure, MAP = mean arterial pressure, Pcc = critical closing pressure, Pmsf = mean systemic filling pressure.

over 10–12 seconds approximates Pcc, following a logarithmic decay pattern (88).

Pmsf Arm

Rapid inflation of a cuff (> 50 mm Hg above SAP) around the upper arm transiently equilibrates arterial and venous pressures, reflecting Pmsf (89, 90). A rapid cuff inflator (0.3 s) (91) or pneumatic tourniquet (1.4 s) (92, 93) should be used. Measuring only arterial pressure until it asymptotes (~18–20 s) is often sufficient (91). This method is applicable in awake patients, those on mechanical ventilation, or with arrhythmias.

Pmsf Analogue

Derived mathematically from the Guyton model ($CO = VR = [Pmsf - CVP]/RVR$) (90, 91, 94–97), where CVP = central venous pressure, RVR = resistance to venous return, and VR = venous return. The Pmsf analogue uses a formula: $Pmsf = a \times CVP + b \times MAP + c \times CO$, where $a + b = 1$, and constants are based on veno-arterial compliance ($a = 0.96$, $b = 0.04$) (94, 96–98). A smartphone app, iGuyton (26), can automate this calculation if CO, MAP, and Pra are known.

Pmsf Inspiratory Hold

This oldest method involves brief end-inspiratory holds (2–10 cm H₂O for 6–10 s), plotting Pra against CO. The zero-flow intercept indicates Pmsf (89, 90). It requires controlled mechanical ventilation without spontaneous breaths or arrhythmias.

ARTERIAL BLOOD PRESSURE AND MACRO-TO-MICROCIRCULATORY COUPLING

Macro-to-microcirculatory coupling (or hemodynamic coherence) refers to the concept where improving systemic macrocirculatory variables, such as MAP and CO, leads to concurrent improvements in microcirculatory flow and reversal of tissue hypoperfusion (99). However, in septic shock, achieving these macrocirculatory targets does not always result in adequate tissue perfusion to meet the metabolic demands of tissues (6). The relationship between increased MAP and microcirculatory perfusion improvement can be

variable. The improvement largely depends on factors such as the balance between systemic TPP reaching a minimal threshold and the existence of a vascular waterfall between the Pcc and the Pmsf, where a decrease in Pcc leads to increased blood flow (100).

Assessing microcirculatory flow at the bedside is challenging. Studies have shown that after increasing MAP to 80–85 mm Hg in hypotensive septic shock patients, microvascular flow responses are heterogeneous. Patients with initially low microvascular flow tended to show improvement at higher MAP levels, whereas those with normal baseline flow may see no improvement, or even a reduction, in flow. In some cases, CRT, a marker of microcirculatory flow, may either increase or decrease when vasopressors are used to raise MAP by the same amount in different patients (101). Additionally, if tissues have entered a state of senescence, simply restoring flow may not improve function. If perfusion pressure is normalized but organ function remains impaired, other factors beyond perfusion pressure are likely responsible for the dysfunction.

The ANDROMEDA-SHOCK trial introduced a test to assess macro-to-microcirculatory coupling by evaluating CRT responses to changes in MAP (102). A positive CRT response to higher MAP indicates that maintaining this higher MAP could be beneficial. Conversely, identifying patients where increasing norepinephrine doses worsen tissue perfusion signals the need to reconsider the MAP target.

Optimizing macrohemodynamics is only the first step in managing septic shock. While MAP is directly related to blood flow, normalizing it does not always improve microcirculatory conditions. MAP values greater than 65 mm Hg generally enhances TPP, which, in turn, may improve tissue perfusion, especially if Pcc is significantly higher than Pmsf. This improvement in MAP can be achieved through fluid resuscitation in fluid-responsive patients. However, if fluid alone does not raise MAP or improve Pcc, vasopressors should be initiated. If MAP exceeds 65 mm Hg and TPP is greater than 35 mm Hg, and the Pcc to Pmsf difference is greater than 10 mm Hg, but hypoperfusion persists, further fluid or vasopressor administration is unlikely to improve perfusion and may instead increase complications. But these final assumptions need to be formally tested before becoming standard practices.

CONCLUSIONS

The relationship between arterial pressure and outcomes in septic shock is complex and heterogeneous. Focusing on critical parameters—such as TPP, the vascular waterfall, and individual treatment effects—enables a more personalized approach, focused on perfusion rather than pressure alone. Persistent hypoperfusion despite adequate macrocirculatory targets marks a pivotal moment when microcirculatory assessment becomes essential to guide therapy and avoid potentially harmful escalation of fluids or vasopressors. Bedside measurements of SAP, MAP, DAP, Pcc, and Pmsf offer practical tools for monitoring and tailoring treatment. Future clinical trials are needed to validate these metrics and refine resuscitation strategies in septic shock.

- 1 Critical Care Medicine, Organ Transplant Center of Excellence, King Faisal Specialist Hospital & Research Centre (KFSHRC), Riyadh, Kingdom of Saudi Arabia.
- 2 Critical Care Institute, Cleveland Clinic, Abu Dhabi, United Arab Emirates.
- 3 Critical Care Medicine, Organ Transplant Center of Excellence, King Faisal Specialist Hospital & Research Centre (KFSHRC), Riyadh, Kingdom of Saudi Arabia.
- 4 College of Medicine, Alfaisal University, Riyadh, Kingdom of Saudi Arabia.
- 5 Departamento de Medicina Intensiva, Facultad de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile.
- 6 Department of Critical Care Medicine, University of Pittsburgh, Pittsburgh, PA.

Dr. Pinsky received funding from Edwards's Lifesciences; he received honoraria for presenting industry-sponsored lectures from Becton-Dickinson. The remaining authors have disclosed that they do not have any potential conflicts of interest.

For information regarding this article, E-mail: dr.carlossanchez@hotmail.com

REFERENCES

1. Singer M, Deutschman CS, Seymour CW, et al: The third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA* 2016; 315:801–810
2. Jarczak D, Kluge S, Nierhaus A: Sepsis—pathophysiology and therapeutic concepts. *Front Med* 2021; 8:628302
3. Varpula M, Tallgren M, Saukkonen K, et al: Hemodynamic variables related to outcome in septic shock. *Intensive Care Med* 2005; 31:1066–1071
4. Ackland GL, Brudney CS, Cecconi M, et al; Perioperative Quality Initiative-3 workgroup: Perioperative quality initiative consensus statement on the physiology of arterial blood pressure control in perioperative medicine. *Br J Anaesth* 2019; 122:542–551
5. Sessler CN, Shepherd W: New concepts in sepsis. *Curr Opin Crit Care* 2002; 8:465–472
6. Bakker J, Ince C: Monitoring coherence between the macro and microcirculation in septic shock. *Curr Opin Crit Care* 2020; 26:267–272
7. De Santis P, De Fazio C, Franchi F, et al: Incoherence between systemic hemodynamic and microcirculatory response to fluid challenge in critically ill patients. *J Clin Med* 2021; 10:507
8. He HW, Long Y, Liu DW, et al: Resuscitation incoherence and dynamic circulation-perfusion coupling in circulatory shock. *Chin Med J (Engl)* 2019; 132:1218–1227
9. Pandey NR, Bian YY, Shou ST: Significance of blood pressure variability in patients with sepsis. *World J Emerg Med* 2014; 5:42–47
10. Hall JE: *Guyton and Hall Textbook of Medical Physiology*. 13th Edition. Philadelphia, PA, W.B.Saunders, 2015
11. Evans L, Rhodes A, Alhazzani W, et al: Surviving sepsis campaign: International guidelines for management of sepsis and septic shock 2021. *Intensive Care Med* 2021; 47:1181–1247
12. Nichols WW, O'Rourke MF: Pulsatile pressure–flow relations. In: *McDonald's Blood Flow in Arteries: Theoretical, Experimental and Clinical Principles*. Chapter 5. Fifth Edition. Nichols WW, O'Rourke MF (Eds). London, Hodder Arnold, 2011, pp 111–129
13. Safar ME, Levy BI, Struijker-Boudier H: Current perspectives on arterial stiffness and pulse pressure in hypertension and cardiovascular diseases. *Circulation* 2003; 107:2864–2869
14. Izzo JL Jr, Shykoff BE: Arterial stiffness: Clinical relevance, measurement, and treatment. *Rev Cardiovasc Med* 2001; 2:29–34, 37
15. Magder S: The meaning of blood pressure. *Crit Care* 2018; 22:257
16. Pinsky MR, García MIM, Dubin A: Significance of critical closing pressures (starling resistors) in arterial circulation. *Crit Care* 2024; 28:127
17. Burton AC: On the physical equilibrium of small blood vessels. *Am J Physiol* 1951; 164:319–329
18. Ostergaard L, Granfeldt A, Secher N, et al: Microcirculatory dysfunction and tissue oxygenation in critical illness. *Acta Anaesthesiol Scand* 2015; 59:1246–1259
19. Chandrasekhar A, Padrós-Valls R, Pallarès-López R, et al: Tissue perfusion pressure enables continuous hemodynamic evaluation and risk prediction in the intensive care unit. *Nat Med* 2023; 29:1998–2006
20. Pinsky MR: The meaning of cardiac output. *Intensive Care Med* 1990; 16:415–417
21. Pinsky MR: A rose by any other name: Cardiac output. *Crit Care Med* 2001; 29:2021–2022
22. Henderson WR, Griesdale DE, Walley KR, et al: Clinical review: Guyton—the role of mean circulatory filling pressure and right atrial pressure in controlling cardiac output. *Crit Care* 2010; 14:243
23. Guyton AC, Polizo D, Armstrong GG: Mean circulatory filling pressure measured immediately after cessation of heart pumping. *Am J Physiol* 1954; 179:261–267

24. Maas JJ, de Wilde RB, Aarts LP, et al: Determination of vascular waterfall phenomenon by bedside measurement of mean systemic filling pressure and critical closing pressure in the intensive care unit. *Anesth Analg* 2012; 114:803–810
25. Rothe CF: Mean circulatory filling pressure: Its meaning and measurement. *J Appl Physiol* 1985; 74:499–509
26. Bertini P: iGuyton (Version 2.1) [Mobile App]. 2018. Available at: <https://apps.apple.com/app/iguyton/id1397046431>. Accessed July 23, 2018
27. Dargent A, Dumargne H, Labruyère M, et al: Role of the interstitium during septic shock: A key to the understanding of fluid dynamics? *J Intensive Care* 2023; 11:44
28. López-Magaña JA, Richards HK, Radolovich DK, et al: Critical closing pressure: Comparison of three methods. *J Cereb Blood Flow Metab* 2009; 29:987–993
29. Sygítowicz G, Sitkiewicz D: Molecular mechanisms of organ damage in sepsis: An overview. *Braz J Infect Dis* 2020; 24:552–560
30. Daering N, Al-Hasan MN: The progression of sepsis from physiologic systemic inflammatory response to immune dysregulation due to life-threatening infections. *Explor Immunol* 2022; 2:665–676
31. Pinsky MR, Guarracino F: How to assess ventriculoarterial coupling in sepsis. *Curr Opin Crit Care* 2020; 26:313–318
32. Monge García MI, Santos A: Understanding ventriculo-arterial coupling. *Ann Transl Med* 2020; 8:795
33. Pinsky MR, Guarracino F: Pathophysiological implications of ventriculoarterial coupling in septic shock. *Intensive Care Med* 2023; 11:87
34. García MI, Romero MG, Cano AG, et al: Dynamic arterial elastance as a predictor of arterial pressure response to fluid administration: A validation study. *Crit Care* 2014; 18:626
35. Bertini P: iElastance (Version 4.1.0) [Mobile App]. 2024. Available at: <https://apps.apple.com/app/ielastance/id556528864> (iOS); <https://play.google.com/store/apps/details?id=unilab.icu.iElastance> (Android). Accessed September 25, 2024
36. Pinsky MR: Protocolized cardiovascular management based on ventricular-arterial coupling. In: *Functional Hemodynamic Monitoring. Update in Intensive Care Medicine*. Pinsky MR, Payen D (Eds). Berlin, Germany, Springer-Verlag, 2006, pp 381–395
37. Monge García MI, Jian Z, Hatib F, et al: Relationship between intraventricular mechanical dyssynchrony and left ventricular systolic and diastolic performance: An in vivo experimental study. *Physiol Rep* 2023; 11:e15607
38. Guinot PG, Bernard E, Levrard M, et al: Dynamic arterial elastance predicts mean arterial pressure decrease associated with decreasing norepinephrine dosage in septic shock. *Crit Care* 2015; 19:14
39. Sanchez JIA, Ruiz SS, Ruiz JDC, et al: Predictive value of dynamic arterial elastance for vasopressor withdrawal: A systematic review and meta-analysis. *Ann Int Care* 2024; 14:108
40. Monge García MI, Jian Z, Hatib F, et al: Dynamic arterial elastance as a ventriculo-arterial coupling index: An experimental animal study. *Front Physiol* 2020; 11:284
41. Alvarado-Sánchez JI, Salazar-Ruiz S, Caicedo-Ruiz JD, et al: Predictive value of dynamic arterial elastance for vasopressor withdrawal: A systematic review and meta-analysis. *Ann Intensive Care* 2024; 14:108
42. Zhou X, Pan W, Chen B, et al: Predictive performance of dynamic arterial elastance for arterial pressure response to fluid expansion in mechanically ventilated hypotensive adults: A systematic review and meta-analysis of observational studies. *Ann Intensive Care* 2021; 11:119
43. Alvarado Sánchez JI, Montañez-Nariño AV, Cárdenas-Bolívar YR, et al: Efficacy and safety of dynamic arterial elastance for weaning vasopressor support in septic shock patients: A randomized controlled trial protocol. *BMJ Open* 2024; 14:e086388
44. London GM, Guerin AP: Influence of the arterial pulse and reflected waves on blood pressure and cardiac function. *Am Heart J* 1999; 138:S220–S224
45. Wilkinson IB, MacCallum H, Flint L, et al: The influence of heart rate on augmentation index and central arterial pressure in humans. *J Physiol* 2000; 525:263–270
46. Hai PD, Phuong LL, Dung NM, et al: Subclinical left ventricular systolic dysfunction in patients with septic shock based on sepsis-3 definition: A speckle-tracking echocardiography study. *Crit Care Res Pract* 2020; 2020:6098654
47. Flint AC, Conell C, Ren X, et al: Effect of systolic and diastolic blood pressure on cardiovascular outcomes. *N Engl J Med* 2019; 381:243–251
48. Tang Y, Sorenson J, Lanspa M, et al: Systolic blood pressure variability in patients with early severe sepsis or septic shock: A prospective cohort study. *BMC Anesthesiology* 2017; 17:82
49. Moran D, Epstein Y, Keren G, et al: Calculation of mean arterial pressure during exercise as a function of heart rate. *Appl Human Sci* 1995; 14:293–295
50. Rodriguez R, Cucci M, Kane S, et al: Novel vasopressors in the treatment of vasodilatory shock: A systematic review of angiotensin II, selexpressin, and terlipressin. *J Intensive Care Med* 2020; 35:327–337
51. Asfar P, Meziani F, Hamel JF, et al; SEPSISPAM Investigators: High versus low blood-pressure target in patients with septic shock. *N Engl J Med* 2014; 370:1583–1593
52. Zhao L, Fan Y, Wang Z, et al: The blood pressure targets in sepsis patients with acute kidney injury: An observational cohort study of multiple ICUs. *Front Immunol* 2022; 13:1060612
53. Malbrain ML, De Waele JJ, De Keulenaer BL: What every ICU clinician needs to know about the cardiovascular effects caused by abdominal hypertension. *Anaesthesiol Intensive Ther* 2015; 47:388–399
54. Cheatham ML: Abdominal compartment syndrome: Pathophysiology and definitions. *Scand J Trauma Resusc Emerg Med* 2009; 17:10
55. Maughan WL, Sunagawa K, Burkhoff D, et al: Effect of heart rate on the canine end-systolic pressure-volume relationship. *Circulation* 1985; 72:654–659
56. Magder S, Guerard B: Heart-lung interactions and pulmonary buffering: Lessons from a computational modeling study. *Respir Physiol Neurobiol* 2012; 182:60–70
57. Monge Garcia MI, Guijo Gonzalez P, Gracia Romero M, et al: Effects of arterial load variations on dynamic arterial elastance: An experimental study. *Br J Anaesth* 2017; 118:938–946

58. Rahman F, Al Rifai M, Blaha MJ, et al: Relation of diastolic blood pressure and coronary artery calcium to coronary events and outcomes (from the multi-ethnic study of atherosclerosis). *Am J Cardiol* 2017; 120:1797–1803
59. Shi R, Hamzaoui O, De Vita N, et al: Vasopressors in septic shock: Which, when, and how much? *Ann Transl Med* 2020; 8:794
60. Hamzaoui O, Shi R: Early norepinephrine use in septic shock. *J Thorac Dis* 2020; 12:S72–S77
61. Xiao W, Liu W, Zhang J, et al: The association of diastolic arterial pressure and heart rate with mortality in septic shock: A retrospective cohort study. *Eur J Med Res* 2022; 27:285
62. Holder AL, Gupta N, Lulaj E, et al: Predictors of early progression to severe sepsis or shock among emergency department patients with nonsevere sepsis. *Int J Emerg Med* 2016; 9:10
63. Gao Z, Li C, Chen H, et al: Association between diastolic blood pressure during the first 24 h and 28-day mortality in patients with septic shock: A retrospective observational study. *Eur J Med Res* 2023; 28:329
64. Kattan E, Bakker J, Estenssoro E, et al: Hemodynamic phenotype-based, capillary refill time-targeted resuscitation in early septic shock: The ANDROMEDA-SHOCK-2 randomized clinical trial study protocol. *Rev Bras Ter Intensiva* 2022; 34:96–106
65. Thompson BT: In septic shock, MAP targets of 80 to 85 mm Hg vs 65 to 70 mm Hg did not differ for 28-day mortality. *Ann Intern Med* 2014; 161:JC7
66. Dilken O, Ergin B, Ince C: Assessment of sublingual microcirculation in critically ill patients: Consensus and debate. *Ann Transl Med* 2020; 8:793
67. Cooper ES, Silverstein DC: Fluid therapy and the microcirculation in health and critical illness. *Front Vet Sci* 2021; 8:625708
68. Collet M, Huot B, Barthelemy R, et al: Influence of systemic hemodynamics on microcirculation during sepsis. *J Crit Care* 2019; 52:213–218
69. Khanna AK, Kinoshita T, Natarajan A, et al: Association of systolic, diastolic, mean, and pulse pressure with morbidity and mortality in septic ICU patients: A nationwide observational study. *Ann Intensive Care* 2023; 13:9
70. Lee SH, Kim YJ, Yu GN, et al: Pulse pressure during the initial resuscitative period in patients with septic shock treated with a protocol-driven resuscitation bundle therapy. *Korean J Intern Med* 2021; 36:924–931
71. Jouffroy R, Gilbert B, Tourtier JP, et al: Prehospital pulse pressure and mortality of septic shock patients cared for by a mobile intensive care unit. *BMC Emerg Med* 2023; 23:97
72. Michard F, Teboul JL: Predicting fluid responsiveness in ICU patients: A critical analysis of the evidence. *Chest* 2002; 121:2000–2008
73. Yang X, Du B: Does pulse pressure variation predict fluid responsiveness in critically ill patients? A systematic review and meta-analysis. *Crit Care* 2014; 18:650
74. Monnet X, Malbrain MLNG, Pinsky MR: The prediction of fluid responsiveness. *Intensive Care Med* 2023; 49:83–86
75. Ospina-Tascón GA, Teboul JL, Hernandez G, et al: Diastolic shock index and clinical outcomes in patients with septic shock. *Ann Intensive Care* 2020; 10:41
76. Kaufmann T, Cox EGM, Wiersema R, et al; SICS Study Group: Non-invasive oscillometric versus invasive arterial blood pressure measurements in critically ill patients: A post hoc analysis of a prospective observational study. *J Crit Care* 2020; 57:118–123
77. Kim SH, Lilot M, Sidhu KS, et al: Accuracy and precision of continuous noninvasive arterial pressure monitoring compared with invasive arterial pressure: A systematic review and meta-analysis. *Anesthesiology* 2014; 120:1080–1097
78. Wax D, Lin HM, Leibowitz AB, et al: Invasive and concomitant noninvasive intraoperative blood pressure monitoring: Observed difference in measurement and associated therapeutic interventions. *Anesthesiology* 2011; 115:973–978
79. Kim WY, Jun JH, Huh JW, et al: Radial to femoral arterial blood pressure differences in septic shock patients receiving high-dose norepinephrine therapy. *Shock* 2013; 40:527–531
80. Wisanusattra H, Khwannimit B: Agreements between mean arterial pressure from radial and femoral artery measurements in refractory shock patients. *Sci Rep* 2022; 12:8825
81. McEniery CM, Cockcroft JR, Roman MJ, et al: Central blood pressure: Current evidence and clinical importance. *Eur Heart J* 2014; 35:1719–1725
82. Herbert A, Cruickshank JK, Laurent S, et al; Reference Values for Arterial Measurements Collaboration: Establishing reference values for central blood pressure and its amplification in a general healthy population and according to cardiovascular risk factors. *Eur Heart J* 2014; 35:3122–3133
83. Chemla D, Millasseau S, Hamzaoui O, et al: New method to estimate central systolic blood pressure from peripheral pressure: A proof of concept and validation study. *Front Cardiovasc Med* 2021; 8:772613
84. Nichols WW, O'Rourke MF, Vlachopoulos C: *McDonald's Blood Flow in Arteries: Theoretical, Experimental and Clinical Principles*. London, United Kingdom, Hodder Arnold, 2011
85. Picone DS, Schultz MG, Otahal P, et al: Accuracy of cuff-measured blood pressure: Systematic reviews and meta-analyses. *J Am Coll Cardiol* 2017; 70:572–586
86. Hatib F, Jansen JR, Pinsky MR: Peripheral vascular decoupling in porcine endotoxemic shock. *J Appl Physiol (1985)* 2011; 111:853–860
87. Andrei S, Bar S, Nguyen M, et al: Effect of norepinephrine on the vascular waterfall and tissue perfusion in vasoplegic hypotensive patients: A prospective, observational, applied physiology study in cardiac surgery. *Intensive Care Med Exp* 2023; 11:52
88. Kottenberg-Assenmacher E, Aleksic I, Eckholt M, et al: Critical closing pressure as the arterial downstream pressure with the heart beating and during circulatory arrest. *Anesthesiology* 2009; 110:370–379
89. Maas JJ, Pinsky MR, Geerts BF, et al: Estimation of mean systemic filling pressure in postoperative cardiac surgery patients with three methods. *Intensive Care Med* 2012; 38:1452–1460
90. Wijnberge M, Sindhunata DP, Pinsky MR, et al: Estimating mean circulatory filling pressure in clinical practice: A systematic review comparing three bedside methods in the critically ill. *Ann Intensive Care* 2018; 8:73
91. Geerts BF, Maas J, de Wilde RB, et al: Arm occlusion pressure is a useful predictor of an increase in cardiac output after fluid

- loading following cardiac surgery. *Eur J Anaesthesiol* 2011; 28:802–806
92. Aya HD, Rhodes A, Fletcher N, et al: Transient stopflow arm arterial-venous equilibrium pressure measurement: Determination of precision of the technique. *J Clin Monit Comput* 2016; 30:55–61
 93. Aya HD, Rhodes A, Chis Ster I, et al: Hemodynamic effect of different doses of fluids for a fluid challenge: A quasi-randomized controlled study. *Crit Care Med* 2017; 45:e161–e168
 94. Parkin WG: Volume state control: A new approach. *Crit Care Resusc* 1999; 1:311–321
 95. Gupta K, Sondergaard S, Parkin G, et al: Applying mean systemic filling pressure to assess the response to fluid boluses in cardiac post-surgical patients. *Intensive Care Med* 2015; 41:265–272
 96. Cecconi M, Aya HD, Geisen M, et al: Changes in the mean systemic filling pressure during a fluid challenge in post-surgical intensive care patients. *Intensive Care Med* 2013; 39:1299–1305
 97. Parkin G, Wright C, Bellomo R, et al: Use of a mean systemic filling pressure analogue during the closed-loop control of fluid replacement in continuous hemodiafiltration. *J Crit Care* 1994; 9:124–133
 98. Crozier TM, Wallace EM, Parkin WG: Haemodynamic assessment in pregnancy and pre-eclampsia: A Guytonian approach. *Pregnancy Hypertens* 2015; 5:177–181
 99. Hernández G, Teboul JL: Is the macrocirculation really dissociated from the microcirculation in septic shock? *Intensive Care Med* 2016; 42:1621–1624
 100. Dubin A, Pozo MO, Casabella CA, et al: Increasing arterial blood pressure with norepinephrine does not improve micro-circulatory blood flow: A prospective study. *Crit Care* 2009; 13:R92
 101. Hernández G, Valenzuela ED, Kattan E, et al: Capillary re-fill time response to a fluid challenge or a vasopressor test: An observational, proof-of-concept study. *Ann Intensive Care* 2024; 14:49
 102. Hernández G, Ospina-Tascón GA, Damiani LP, et al; The ANDROMEDA SHOCK Investigators and the Latin America Intensive Care Network (LIVEN): Effect of a resuscitation strategy targeting peripheral perfusion status vs serum lactate levels on 28-day mortality among patients with septic shock: The ANDROMEDA-SHOCK randomized clinical trial. *JAMA* 2019; 321:654–664